

NOTES

Mutational Evidence for Identity of Penicillin-Binding Protein 5 in *Escherichia coli* with the Major D-Alanine Carboxypeptidase IA Activity

MICHIO MATSUHASHI,¹* SHIGEO TAMAKI,¹ SUSAN J. CURTIS,² AND JACK L. STROMINGER²

Institute of Applied Microbiology, University of Tokyo, Bunkyo-ku, Tokyo 113, Japan,¹ and Harvard University, The Biological Laboratories, Cambridge, Massachusetts 02138²

Received for publication 11 October 1978

The defect in D-alanine carboxypeptidase IA activity in the *dacA*11191 mutant of *Escherichia coli* was correlated with a defect in the release of penicillin G from penicillin-binding protein 5. The results suggest that penicillin-binding protein 5 catalyzes the major D-alanine carboxypeptidase IA activity of the wild type and that the mutation results in a defect in the deacylation step catalyzed by this enzyme.

Purified D-alanine carboxypeptidase IA in *Escherichia coli* consists of two polypeptides which can be separated by sodium dodecyl sulfate-acrylamide gel electrophoresis (12). Both proteins bind penicillin G and have been identified as penicillin-binding proteins (PBP) 5 and 6 of *E. coli* (10; see also 9). Both proteins also release the bound penicillin G (8); these abilities represent weak β -lactamase activities (12). Tamura et al. (12) and Curtis and Strominger (1) demonstrated that both activities, i.e., D-alanine carboxypeptidase and release of enzyme-bound penicillin G, are sensitive to sulfhydryl reagents such as *p*-chloromercuribenzoate (pCMB). These reagents have no effect on penicillin G binding and, correspondingly, pCMB causes accumulation of an acyl enzyme intermediate of the carboxypeptidase reaction derived from a synthetic substrate, diacetyl-L-lysyl-D-alanyl-D-alanine (1). The same intermediate was subsequently obtained in higher yield with the more reactive ester substrate diacetyl-L-lysyl-D-alanyl-D-lactate (7). Thus, it was proposed (1, 12) that the binding of penicillin G to D-alanine carboxypeptidase IA and its release correspond to substeps of the total D-alanine carboxypeptidase reaction. The first step (step 1), which is rather insensitive to pCMB, is the formation of an acyl enzyme intermediate with concomitant release of the terminal D-alanine of a substrate such as diacetyl-L-lysyl-D-alanyl-D-alanine or UDP-*N*-acetylmuramyl-pentapeptide. The second step (step 2), which is sensitive to pCMB, is the transfer of the acyl group to water. If the substrate is penicillin, the sum of steps 1 and 2

is the β -lactamase reaction. This paper describes mutational evidence for the identity of these two enzymatic reactions and also suggests the possibility that only PBP 5 is responsible for the enzymatic activity, D-alanine carboxypeptidase IA.

The *dacA*11191 mutant of *E. coli* strain JE11191 (*F*⁻ *thr leu trp his thyA argH thi lacY malA mtl mel tonA rpsL dacA*), which lacks D-alanine carboxypeptidase IA activity, was described previously (4). Isogenic strains containing either the *dacA*⁺ or the *dacA* allele were obtained by transduction of strain TMRL-122 (*F*⁻ *proA purB his thi lacY galK rpsL leuS dacB*) with phage P-1 grown in the *leuS*⁺ *dacA*11191 strain. *leuS*⁺ transductants were selected. Strain TMRL-122 was obtained by *argG*⁺ transduction of strain TMRL-12 (*F*⁻ *proA purB his argG thi lacY galK rpsL leuS*) with phage P-1 grown in an *argG*⁺ *dacB* strain that lacks D-alanine carboxypeptidase IB activity (5). The presence of the *dacB* mutation greatly facilitates the determination of D-alanine carboxypeptidase IA activity. The map positions of the *dacA* and *dacB* genes and the method used for preparation of strains with *dacA dacB* double mutations will be described elsewhere (4, 10a; Y. Hirota, Y. Nishimura, Y. Takagaki, I. N. Maruyama, and M. Matsuhashi, in preparation).

The defect of D-alanine carboxypeptidase activities in these *dacA dacB* strains was demonstrated by the two different assay methods previously described (Table 1): experiment 1, the exchange reaction of [¹⁴C]glycine with terminal D-alanine of UDP-MurNAc-pentapeptide (L-

TABLE 1. Enzyme activities in isogenic *dacA⁺ dacB* and *dacA dacB* strains^a

Expt	Strain and genotype	Addition	Enzyme activity (pmol/mg of protein per h)
1	TMRL-1224 (<i>dacA⁺ dacB</i>)	None	171
		0.1 mM pCMB	19
		0.1 mM pCMB plus 1 mM 2-mercaptoethanol	166
	TMRL-1222 (<i>dacA dacB</i>)	None	1
2	TMRL-1224 (<i>dacA⁺ dacB</i>)	None	826
		3 µg of penicillin G per ml	540
		10 mM MgCl ₂	470
	TMRL-1222 (<i>dacA dacB</i>)	None	12

^a D-Alanine carboxypeptidase activity was assayed in an extract [0.05 M tris(hydroxymethyl)amino-methane (Tris)-hydrochloride buffer, pH 7.5] of a sonic lysate of late-log-phase cells that had been cultured in L-broth (3) supplemented with 20 µg of thymine per ml. For [¹⁴C]glycine exchange (pseudotranspeptidase reaction, experiment 1), the reaction mixture contained in a final volume of 30 µl: 2 µmol of Tris-hydrochloride buffer, pH 9.0; 0.1 µmol of UDP-MurNAc-L-Ala-D-Glu-m-A₂pm-D-Ala-D-Ala; 1 nmol of [¹⁴C]glycine (102 µCi/µmol, New England Nuclear); sonic cell extract (50 to 100 µg of protein); 0.03 µmol of 2-mercaptoethanol; and 0.67% (wt/vol) Triton X-100. It was incubated for 60 min at 30°C, heated at 100°C for 1 min, and chromatographed on Whatman no. 3 MM filter paper in the solvent isobutyric acid-1 M ammonia (5:3, vol/vol). After chromatography, the radioactive area on the paper was detected in a spark chamber (Birchover Co., England), and radioactivity on the paper corresponding to the product was counted in a liquid scintillation spectrometer by immersing the paper in toluene-PPO (2,5-diphenyloxazole)-POPOP [2,2'-p-phenylene-bis-(5-phenyloxazole)] (1 liter:4 g:100 mg). For assay of D-[¹⁴C]alanine release (experiment 2), the reaction mixture contained, in a final volume of 30 µl: 3 µmol of Tris-hydrochloride buffer (pH 9.0); 0.3 nmol of UDP-MurNAc-L-Ala-D-Glu-m-A₂pm-D-[¹⁴C]Ala-D-[¹⁴C]Ala (20 µCi/µmol, prepared as described previously [2]); sonic cell extract (50 to 100 µg of protein); 0.05 µmol of 2-mercaptoethanol; and 1% (wt/vol, final) Triton X-100. The reaction was carried out for 60 min at 30°C. Subsequent procedures were as described above. Results on other isogenic strains (TMRL-12211 and TMRL-1228, etc.) were similar.

Ala-D-Glu-m-A₂pm-D-Ala-D-Ala) (4-6, 12); and experiment 2, release of terminal D-[¹⁴C]alanine from UDP-MurNAc-pentapeptide labeled in the terminal D-alanyl-D-alanine group, D-[¹⁴C]Ala-D-[¹⁴C]Ala (2, 4, 5). Strains *dacA dacB* showed scarcely any activity in either enzyme assay (Table 1). *dacA⁺ dacB* strains showed apprecia-

ble activity in both reactions, and this activity was due to D-alanine carboxypeptidase IA activity, as indicated by its intermediate sensitivity to penicillin G, slight inhibition by magnesium ion, inhibition by 100 µM pCMB, and protection from pCMB inhibition by excess 2-mercaptoethanol.

The isogenic *dacA⁺* and *dacA* strains (both containing the *dacB* mutation) gave similar patterns of PBPs. The differences in the amounts of PBP 5 and PBP 6 in the *dacA* mutant strain JE11191 compared to the amounts in the parent strain PA3092 observed previously (4) were not observed in the isogenic *dacA⁺* and *dacA* strains. Therefore, the changes in the amounts of PBP 5 and 6 are not due to the *dacA* mutation itself, but rather to some other genetic or physiological changes occurring in the mutant JE11191.

Curtis and Strominger previously suggested (1; see also 4) that the *dacA11191* mutation may produce a defect in carboxypeptidase IA similar to the alteration produced by sulfhydryl reagents, which inhibits step 2 of the enzyme reaction. Thus, release of [¹⁴C]penicillin G from PBP 5 and 6 was examined in isogenic *dacA⁺ dacB* and *dacA dacB* strains. The release of [¹⁴C]penicillin G from PBP 5 was almost completely blocked in *dacA* strains, whereas the release from PBP 6, which is normally much slower than that from wild-type PBP 5, was unchanged in *dacA* strains (Fig. 1). The results were confirmed with 12 *dacA⁺ dacB* and 14 *dacA dacB* strains and with 7 *dacA⁺ dacB⁺* and 5 *dacA dacB⁺* strains (total, 38 strains). The 12 strains containing the *dacB⁺* gene consisted of 10 isogenic strains isolated by a method similar to that described above, the original *dacA* mutant strain JE11191, and its *dacA⁺* parent strain, PA3092.

The following conclusions are drawn from these results: (i) the *dacA11191* mutation, located at 13.7 min on the *E. coli* chromosome, causes a defect in the release of bound penicillin G from PBP 5; (ii) PBP 5 may be responsible for the major enzyme activity of D-alanine carboxypeptidase IA; and (iii) the penicillin G-releasing activity of PBP 5 appears to be identical to step 2 of the D-alanine carboxypeptidase reaction, in which the acyl enzyme intermediate derived from substrate (the terminal D-alanine having been removed) is transferred to water or another acceptor.

Unlike PBP 5, PBP 6 does not seem to be responsible for the major D-alanine carboxypeptidase reaction, because this protein in *dacA11191* mutant cells can bind [¹⁴C]penicillin G and release it even though the crude cell extract of the mutants lacks D-alanine carboxy-




FIG. 1. Defect in the release of [14 C]penicillin G from PBP 5 in the *dacA*11191 mutant. Release of penicillin from PBP 5 and PBP 6 in the isogenic strains (A) TMRL-12211 (*dacA*⁺ *dacB*) and (B) TMRL-1228 (*dacA* *dacB*) is shown. Experimental procedures were essentially as described previously (8–11). Cells in the log phase (about 10^8 cells per ml), grown at 30°C in L-broth (3) supplemented with 20 μ g of thymine per ml, were sonically treated in 0.01 M sodium phosphate buffer, pH 7.0, and the membrane fraction was obtained by fractional centrifugation between $3,000 \times g$ (10 min) and $100,000 \times g$ (30 min). The fraction was washed with the same buffer and suspended in 0.05 M sodium phosphate buffer, pH 7.0 (protein concentration, 20 mg/ml). For binding of [14 C]penicillin G, a mixture of 30 μ l of membrane suspension and 3 μ l of [14 C]penicillin G (50 μ Ci/ μ mol, Radiochemical Centre, Amersham, England; 1 mM) was incubated for 10 min at 30°C. After binding of [14 C]penicillin G to the proteins as described above, 0.9 mg of unlabeled penicillin G (Takeda Chemical Industry Co., Osaka; 800-fold excess) was added to the mixture, and the incubation was continued at 30°C for the indicated periods. Then the reaction mixture was mixed with 2 μ l of a solution containing 90 μ g of unlabeled penicillin G and 15% (wt/vol) Sarkosyl (Ciba Geigy). The Sarkosyl-insoluble outer membrane fraction was removed by centrifugation at $12,000 \times g$ for 30 min at 20°C, and the supernatant was mixed with 20 μ l of a mixture containing 2.25% (wt/vol) sodium dodecyl sulfate, 0.15 M tris(hydroxymethyl)aminomethane - hydrochloride buffer (pH 6.8), 22.5% (wt/vol) glycerol, 0.0015% (wt/vol) bromophenol blue, and 25% (wt/vol) 2-mercaptoethanol and heated for 2 min at 100°C. Then it was subjected to sodium dodecyl sulfate-polyacrylamide slab gel electrophoresis. A separating gel containing 7.5% (wt/vol) acrylamide with 1.3% (wt/wt) cross-linking was used (11). The fluorogram of the slab gel was prepared as described previously (8, 11). Results on other isogenic strains (e.g., TMRL-1224 and TMRL-1222) were similar.

peptidase activity. Further work involving separation of PBP 6 free from PBP 5 is, however, necessary for the identification of the enzymatic activity of PBP 6.

It is possible that the gene for D-alanine car-

boxypeptidase IA (*dacA*) and the gene for PBP 5 are located very close together on the *E. coli* chromosome and that the first *dacA* mutant isolated, strain JE11191, had independent mutations of both of these genes. It seems very unlikely that the two properties of *dacA* are due to a double mutation, since 36 transductants, 18 of which were *dacA*⁺ and 18 of which were *dacA*, were isolated. These transductants were selected for the *leuS*⁺ marker, which is 90% linked to the *dacA* marker by cotransduction with phage P-1. Therefore, the finding that in each of the 36 transductants either both activities were defective or both were normal indicates that these two enzymatic defects are due to a single mutation. The recent observation of Tamaki et al. (10a) that the *dacA* mutation results in super-sensitivity of *E. coli* cells to various β -lactam antibiotics indicates that the *dacA*⁺ revertant can be obtained from *dacA* cells by selecting cells that are less sensitive to β -lactam antibiotics.

This research was supported by a Public Health Service research grant from the National Institute of Allergy and Infectious Diseases (AI 09152) and Ministry of Education, Japan.

LITERATURE CITED

1. Curtis, S. J., and J. L. Strominger. 1978. Effects of sulphydryl reagents on the binding and release of penicillin G by D-alanine carboxypeptidase IA of *Escherichia coli*. J. Biol. Chem. 253:2584–2588.
2. Izaki, K., M. Matsuhashi, and J. L. Strominger. 1966. Glycopeptide transpeptidase and D-alanine carboxypeptidase: penicillin-sensitive enzymatic reactions. Proc. Natl. Acad. Sci. U.S.A. 55:656–663.
3. Lennox, E. S. 1955. Transduction of linked genetic characters of the host by bacteriophage P1. Virology 1: 190–206.
4. Matsuhashi, M., I. N. Maruyama, Y. Takagaki, S. Tamaki, Y. Nishimura, and Y. Hirota. 1978. Isolation of a mutant of *Escherichia coli* lacking penicillin-sensitive D-alanine carboxypeptidase IA. Proc. Natl. Acad. Sci. U.S.A. 75:2631–2635.
5. Matsuhashi, M., Y. Takagaki, I. N. Maruyama, S. Tamaki, Y. Nishimura, H. Suzuki, U. Ogino, and Y. Hirota. 1977. Mutants of *Escherichia coli* lacking in highly penicillin-sensitive D-alanine carboxypeptidase activity. Proc. Natl. Acad. Sci. U.S.A. 74: 2976–2979.
6. Pollock, J. J., M. Nguyen-Disteche, J. M. Ghuyssen, J. Coyette, R. Linder, M. R. J. Salton, K. S. Kim, H. R. Perkins, and P. Reynolds. 1974. Fractionation of the DD-carboxypeptidase-transpeptidase activities solubilized from membranes of *Escherichia coli* K12, strain 44. Eur. J. Biochem. 41:439–446.
7. Rasmussen, J. R., and J. L. Strominger. 1978. Utilization of a depsipeptide substrate for trapping acyl-enzyme intermediates of penicillin-sensitive D-alanine carboxypeptidases. Proc. Natl. Acad. Sci. U.S.A. 75: 84–88.
8. Spratt, B. G. 1977. Properties of the penicillin-binding proteins of *Escherichia coli* K12. Eur. J. Biochem. 72: 341–352.
9. Spratt, B. G., and A. B. Pardee. 1975. Penicillin-binding

- proteins and cell shape in *E. coli*. *Nature* (London) **254**: 516-517.
10. **Spratt, B. G., and J. L. Strominger.** 1976. Identification of the major penicillin-binding proteins of *Escherichia coli* as D-alanine carboxypeptidase IA. *J. Bacteriol.* **127**: 660-663.
- 10a. **Tamaki, S., J. Nakagawa, I. N. Maruyama, and M. Matsuhashi.** 1978. Supersensitivity to β -lactam antibiotics in *Escherichia coli* caused by D-alanine carboxypeptidase IA mutation. *Agric. Biol. Chem.* **42**:2147-2150.
11. **Tamaki, S., S. Nakajima, and M. Matsuhashi.** 1977. Thermosensitive mutation in *Escherichia coli* simultaneously causing defects in penicillin-binding protein-IBs and in enzyme activity for peptidoglycan synthesis *in vitro*. *Proc. Natl. Acad. Sci. U.S.A.* **74**:5472-5476.
12. **Tamura, T., Y. Imae, and J. L. Strominger.** 1976. Purification to homogeneity and properties of two D-alanine carboxypeptidases I from *Escherichia coli*. *J. Biol. Chem.* **251**:414-423.